






# ABEMACICLIB (LY2835219)

## CDK4/6 INHIBITOR

**The safety and efficacy of the agents for uses under investigation have not been established. Pipeline molecules may not receive regulatory approval and become commercially available for the uses being investigated. The information provided about new molecules being studied is for scientific information exchange purposes only with no commercial intent. For more information on our pipeline, please visit [lillyoxooncologypipeline.com](https://lillyoxooncologypipeline.com).**

**This document was commissioned by Lilly Medical and is intended to be used by HCPs for medical, scientific, and educational purposes.**

# ABEMACICLIB CDK4/6 INHIBITOR (LY2835219) | MECHANISM OF ACTION<sup>1</sup>



Shapiro CI, et al<sup>1</sup>

**Reference:** 1. Shapiro GI. *J Clin Oncol.* 2006;24(11):1770-1783.

The safety and efficacy of the agents for uses under investigation have not been established. Pipeline molecules may not receive regulatory approval and become commercially available for the uses being investigated. The information provided about new molecules being studied is for scientific information exchange purposes only with no commercial intent. For more information on our pipeline, please visit [lillyoxooncologypipeline.com](http://lillyoxooncologypipeline.com).

This document was commissioned by Lilly Medical and is intended to be used by HCPs for medical, scientific, and educational purposes.



# ABEMACICLIB CDK4/6 INHIBITOR (LY2835219)

## TARGET

Many human tumors acquire alterations which can lead to the activation of cyclin-dependent kinases (CDKs). These alterations include mutations that directly activate CDK4/6; gene amplifications, which increase expression of various protein activators such as D-type cyclins; as well as genetic losses, which reduce expression of protein inhibitors such as p16. These various mechanisms as well as loss of retinoblastoma (Rb) can lead to an enhanced proliferative potential by decreasing dependency on external growth factors and mitogenic signaling pathways, which are required to stimulate growth under normal conditions.<sup>1,2</sup>

## MOLECULE

Abemaciclib has been shown *in vitro* to be a selective ATP-competitive inhibitor of CDK4/6 kinase activity that prevents the phosphorylation and subsequent inactivation of the Rb tumor suppressor protein, thereby inducing G1 cell-cycle arrest and inhibition of cell proliferation.<sup>3,4</sup>

## CLINICAL DEVELOPMENT

Abemaciclib is being investigated in clinical trials in patients with breast cancer, non-small cell lung cancer, pediatric cancers, or sarcoma.

**References:** 1. Kim JK, Diehl JA. *J Cell Physiol.* 2009;220(2):292-296. 2. Choi YJ, Anders L. *Oncogene.* 2014;33(15):1890-1903. 3. Dempsey JA, et al. AACR Annual Meeting; April 6-10, 2013; Washington, DC. Abstract LB122. 4. Gelbert LM, et al. *Invest New Drugs.* 2014;32(5):825-837.

**The safety and efficacy of the agents for uses under investigation have not been established. Pipeline molecules may not receive regulatory approval and become commercially available for the uses being investigated. The information provided about new molecules being studied is for scientific information exchange purposes only with no commercial intent. For more information on our pipeline, please visit [lillyoxooncologypipeline.com](http://lillyoxooncologypipeline.com).**


**This document was commissioned by Lilly Medical and is intended to be used by HCPs for medical, scientific, and educational purposes.**



# ABEMACICLIB CDK4/6 INHIBITOR (LY2835219)

## CAMPFIRE

A Randomized, Open-Label, Phase 2 Study Evaluating Abemaciclib in Combination With Irinotecan and Temozolomide in Participants With Relapsed or Refractory Ewing's Sarcoma<sup>a</sup>



<sup>a</sup>This clinical trial is being conducted globally. <sup>b</sup>Abemaciclib is administered PO. <sup>c</sup>Irinotecan is administered intravenously. <sup>d</sup>Temozolomide is administered. **Abbreviations:** PO=orally.

Please visit [clinicaltrials.gov](https://clinicaltrials.gov) for more information on this clinical trial [NCT05440786].

The safety and efficacy of the agents for uses under investigation have not been established. Pipeline molecules may not receive regulatory approval and become commercially available for the uses being investigated. The information provided about new molecules being studied is for scientific information exchange purposes only with no commercial intent. For more information on our pipeline, please visit [lillyoxooncologypipeline.com](https://lillyoxooncologypipeline.com).

This document was commissioned by Lilly Medical and is intended to be used by HCPs for medical, scientific, and educational purposes.

# ABEMACICLIB CDK4/6 INHIBITOR (LY2835219)

## CAMPFIRE

**A Randomized, Open-Label, Phase 2 Study Evaluating Abemaciclib in Combination With Irinotecan and Temozolomide in Participants With Relapsed or Refractory Ewing's Sarcoma (cont.)**

## KEY INCLUSION CRITERIA

- Ages 1 to 39
- Ewing's sarcoma or Ewing's sarcoma-like tumor
  - The original pathological report is required; repeat biopsy at progression is not required
- Refractory disease or confirmed radiological progression or recurrence following first or later line of treatment of Ewing's sarcoma or Ewing's sarcoma-like tumor
  - Participants must have one measurable or evaluable lesion as defined by Response Evaluation Criteria in Solid Tumors version 1.1 (RECIST v1.1)
- Lansky score  $\geq 50$  for participants  $< 16$  years of age, and Karnofsky score  $\geq 50$  for participants  $\geq 16$  years of age
- Participants must have discontinued all previous treatments for cancer or investigational agents  $\geq 7$  days after the last dose and must have recovered from the acute effects
- Adequate hematologic and organ function  $\leq 14$  days prior to day 1 of cycle 1
  - Platelets  $\geq 75,000$ /cubic millimeter ( $\text{mm}^3$ ), hemoglobin  $\geq 8$  g/dL, and absolute neutrophil count  $\geq 1000$ /microliter ( $\mu\text{L}$ )
  - Aspartate aminotransferase (AST) and alanine aminotransferase (ALT)  $\leq 3.0 \times$  upper limit of normal (ULN); total bilirubin  $\leq 1.5 \times$  ULN
  - Creatinine clearance or calculated glomerular filtration rate (GFR)  $\geq 60$  mL/min/ $\text{m}^2$  or serum creatinine based on age/gender
- Female participants of childbearing potential must have a negative urine or serum pregnancy test
- Body weight  $\geq 10$  kg
- Must be able to swallow and/or have a gastric/nasogastric tube
  - Participants in the European Union must be able to swallow intact capsules
- Stable or decreasing dose of steroids at least 7 days prior to enrollment
- Life expectancy of at least 8 weeks and able to complete at least 1 cycle of treatment
- Participants/caregivers are able and willing to make themselves available for the duration of the study and are willing to follow study procedures, including adherence to the pharmacokinetic (PK) sampling schedule

## KEY EXCLUSION CRITERIA

- Severe and/or uncontrolled concurrent medical disease or psychiatric illness/social situation that, in the judgment of the investigator, could cause unacceptable safety risks or compromise compliance with the protocol
- Active fungal, bacterial, and/or known severe viral infection, including but not limited to HIV or viral hepatitis A, B, or C
- Prior allogeneic bone marrow or solid organ transplant
- Major surgical procedure, laparoscopic procedure, or significant traumatic injury within 28 days prior to enrollment. Surgical or other wounds must be adequately healed prior to enrollment
- Pregnant or breastfeeding
- Prior treatment with a cyclin-dependent kinase 4 and 6 (CDK4/6) inhibitor
- Progression during prior treatment with irinotecan and/or temozolomide
- Known intolerance or hypersensitivity to any of the study treatments or dacarbazine
- Diagnosed and/or treated for an additional malignancy within 3 years prior to enrollment

Please visit [clinicaltrials.gov](https://clinicaltrials.gov) for more information on this clinical trial [NCT05440786].

The safety and efficacy of the agents for uses under investigation have not been established. Pipeline molecules may not receive regulatory approval and become commercially available for the uses being investigated. The information provided about new molecules being studied is for scientific information exchange purposes only with no commercial intent. For more information on our pipeline, please visit [lillyoxooncologypipeline.com](https://lillyoxooncologypipeline.com).

This document was commissioned by Lilly Medical and is intended to be used by HCPs for medical, scientific, and educational purposes.



## ACTIVE TRIALS CURRENTLY NOT ENROLLING

### **[NCT02107703] Breast Cancer**

MONARCH 2: A Study of Abemaciclib (LY2835219) Combined With Fulvestrant in Women With Hormone-Receptor-Positive, HER2-Negative Breast Cancer

### **[NCT02246621] Breast Cancer**

MONARCH 3: A Study of Nonsteroidal Aromatase Inhibitors Plus Abemaciclib (LY2835219) in Postmenopausal Women With Breast Cancer

### **[NCT03155997] Breast Cancer**

monarchE: Endocrine Therapy With or Without Abemaciclib (LY2835219) Following Surgery in Participants With Breast Cancer

### **[NCT05169567] Breast Cancer**

postMONARCH: Abemaciclib (LY2835219) Plus Fulvestrant Compared to Placebo Plus Fulvestrant in Previously Treated Breast Cancer

### **[NCT04238819] Solid Tumors**

A Study of Abemaciclib (LY2835219) in Combination With Other Anti-Cancer Treatments in Children and Young Adult Participants With Solid Tumors, Including Neuroblastoma

### **[NCT02057133] Breast Cancer**

A Study of LY2835219 (Abemaciclib) in Combination With Therapies for Breast Cancer That Has Spread

### **[NCT02152631] Lung Cancer**

JUNIPER: A Study of Abemaciclib (LY2835219) in Participants With Previously Treated KRAS Mutated Lung Cancer

### **[NCT05288166] Prostate Cancer**

CYCLONE 3: A Study of Abemaciclib (LY2835219) With Abiraterone in Men With Prostate Cancer That Has Spread to Other Parts of the Body and is Expected to Respond to Hormonal Treatment (Metastatic Hormone-Sensitive Prostate Cancer)

**The safety and efficacy of the agents for uses under investigation have not been established. Pipeline molecules may not receive regulatory approval and become commercially available for the uses being investigated. The information provided about new molecules being studied is for scientific information exchange purposes only with no commercial intent. For more information on our pipeline, please visit [lillyoxooncologypipeline.com](https://lillyoxooncologypipeline.com).**

**This document was commissioned by Lilly Medical and is intended to be used by HCPs for medical, scientific, and educational purposes.**

## ACTIVE TRIALS CURRENTLY NOT ENROLLING (CONT.)

### **[NCT02779751] Lung Cancer or Breast Cancer**

A Study of Abemaciclib (LY2835219) in Participants With Non-Small Cell Lung Cancer or Breast Cancer

### **[NCT04975308] Breast Cancer**

EMBER-3: A Study of Imlunestrant, Investigator's Choice of Endocrine Therapy, and Imlunestrant Plus Abemaciclib in Participants With ER+, HER2- Advanced Breast Cancer

### **[NCT02747004] Breast Cancer**

Next MONARCH 1: A Study of Abemaciclib (LY2835219) Plus Tamoxifen or Abemaciclib Alone in Women With Metastatic Breast Cancer

### **[NCT05999968] Prostate Cancer**

Abemaciclib Plus Darolutamide in Prostate Cancer That Has Spread After Initial Treatment

### **[NCT03706365] Prostate Cancer**

CYCLONE 2: A Study of Abiraterone Acetate Plus Prednisone With or Without Abemaciclib (LY2835219) in Participants With Prostate Cancer

### **[NCT02763566] Breast Cancer**

MONARCH plus: A Study of Abemaciclib (LY2835219) in Participants With Breast Cancer

### **[NCT02791334] Solid Tumors**

PACT: A Study of Anti-PD-L1 Checkpoint Antibody (LY3300054) Alone and in Combination in Participants With Advanced Refractory Solid Tumors

### **[NCT04188548] Breast Cancer or Endometrial Cancer**

EMBER: A Study of LY3484356 in Participants With Advanced or Metastatic Breast Cancer or Endometrial Cancer

**The safety and efficacy of the agents for uses under investigation have not been established. Pipeline molecules may not receive regulatory approval and become commercially available for the uses being investigated. The information provided about new molecules being studied is for scientific information exchange purposes only with no commercial intent. For more information on our pipeline, please visit [lillyoxooncologypipeline.com](https://lillyoxooncologypipeline.com).**

**This document was commissioned by Lilly Medical and is intended to be used by HCPs for medical, scientific, and educational purposes.**

Pipeline information is current through July 27, 2024.